

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:SSPTADKO1625

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	JAN 02	STN pricing information for 2008 now available
NEWS	3	JAN 16	CAS patent coverage enhanced to include exemplified prophetic substances
NEWS	4	JAN 28	USPATFULL, USPAT2, and USPATOLD enhanced with new custom IPC display formats
NEWS	5	JAN 28	MARPAT searching enhanced
NEWS	6	JAN 28	USGENE now provides USPTO sequence data within 3 days of publication
NEWS	7	JAN 28	TOXCENTER enhanced with reloaded MEDLINE segment
NEWS	8	JAN 28	MEDLINE and LMEEDLINE reloaded with enhancements
NEWS	9	FEB 08	STN Express, Version 8.3, now available
NEWS	10	FEB 20	PCI now available as a replacement to DPIC
NEWS	11	FEB 25	IFIREF reloaded with enhancements
NEWS	12	FEB 25	IMSPRODUCT reloaded with enhancements
NEWS	13	FEB 29	WPINDEX/WPIDS/WPIX enhanced with ECLA and current U.S. National Patent Classification
NEWS	14	MAR 31	IFICDB, IFIPAT, and IFIUDB enhanced with new custom IPC display formats
NEWS	15	MAR 31	CAS REGISTRY enhanced with additional experimental spectra
NEWS	16	MAR 31	CA/CAPLUS and CASREACT patent number format for U.S. applications updated
NEWS	17	MAR 31	LPCI now available as a replacement to LDPCI
NEWS	18	MAR 31	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS	19	APR 04	STN AnaVist, Version 1, to be discontinued
NEWS	20	APR 15	WPIDS, WPINDEX, and WPIX enhanced with new predefined hit display formats
NEWS	21	APR 28	EMBASE Controlled Term thesaurus enhanced
NEWS	22	APR 28	IMSRSEARCH reloaded with enhancements
NEWS EXPRESS	FEBRUARY 08 CURRENT WINDOWS VERSION IS V8.3, AND CURRENT DISCOVER FILE IS DATED 20 FEBRUARY 2008		
NEWS HOURS	STN Operating Hours Plus Help Desk Availability		
NEWS LOGIN	Welcome Banner and News Items		
NEWS IPC8	For general information regarding STN implementation of IPC 8		

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 13:04:25 ON 13 MAY 2008

=> file casreact

FILE 'CASREACT' ENTERED AT 13:05:03 ON 13 MAY 2008

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 10 May 2008 VOL 148 ISS 20

New CAS Information Use Policies, enter HELP USAGETERMS for details.

\*\*\*\*\*  
\*  
\* CASREACT now has more than 13.8 million reactions \*  
\*  
\*\*\*\*\*

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=>


Uploading C:\Program Files\Stnexp\Queries\10565981.str

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 13:05:38 FILE 'CASREACT'

SCREENING COMPLETE -

0 REACTIONS TO VERIFY FROM

0 DOCUMENTS

100.0% DONE

0 VERIFIED

0 HIT RXNS

0 DOCS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE \*\*COMPLETE\*\*  
BATCH \*\*COMPLETE\*\*

PROJECTED VERIFICATIONS: 0 TO 0

PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1 ( 0 REACTIONS)

=> s l1 full

FULL SEARCH INITIATED 13:06:07 FILE 'CASREACT'

SCREENING COMPLETE - 247 REACTIONS TO VERIFY FROM 33 DOCUMENTS

100.0% DONE 247 VERIFIED 80 HIT RXNS 10 DOCS

SEARCH TIME: 00.00.01

L3 10 SEA SSS FUL L1 ( 80 REACTIONS)

=> d help

The following are valid formats:

ABS ----- GI and AB  
ALL ----- BIB, AB, IND, RE, Single-step Reactions  
APPS ----- AI, PRAI  
BIB ----- AN, plus Bibliographic Data  
CAN ----- List of CA abstract numbers without answer numbers  
CBIB ----- AN, plus Compressed Bibliographic Data  
DALL ----- ALL, delimited (end of each field identified)  
IABS ----- ABS, indented with text labels  
IALL ----- ALL, indented with text labels  
IBIB ----- BIB, indented with text labels  
IND ----- Indexing data  
IPC ----- International Patent Classifications  
ISTD ----- STD, indented with text labels  
OBIB ----- AN, plus Bibliographic Data (original)  
OIBIB ----- OBIB, indented with text labels  
  
SBIB ----- BIB, no citations  
SIBIB ----- IBIB, no citations  
  
MAX ----- Same as ALL  
PATS ----- PI, SO  
SCAN ----- TI and FCRD (random display, no answer number. SCAN  
must be entered on the same line as DISPLAY, e.g.,  
D SCAN.)  
SSRX ----- Single-Step Reactions (Map, Diagram, and Summary for  
all single-step reactions)  
STD ----- BIB, IPC, and NCL  
  
CRD ----- Compact Display of All Hit Reactions  
CRDREF ----- Compact Reaction Display and SO, PY for Reference  
FHIT ----- Reaction Map, Diagram, and Summary for first  
hit reaction  
FHITCBIB --- FHIT, AN plus CBIB  
FCRD ----- First hit in Compact Reaction Display (CRD) format  
FCRDREF ---- First hit in Compact Reaction Display (CRD) format with  
CA reference information (SO, PY). (Default)  
FPATH ----- PATH, plus Reaction Summary for the "long path"  
FSPATH ----- SPATH, plus Reaction Summary for the "short path"  
HIT ----- Reaction Map, Reaction Diagram, and Reaction  
Summary for all hit reactions and fields containing

hit terms

OCC ----- All hit fields and the number of occurrences of the hit terms in each field. Includes total number of HIT, PATH, SPATH reactions. Labels reactions that have incomplete verifications.

PATH ----- Reaction Map and Reaction Diagram for the "long path". Displays all hit reactions, except those whose steps are totally included within another hit reaction which is displayed

RX ----- Hit Reactions (Map, Diagram, Summary for all hit reactions)

RXG ----- Hit Reaction Graphics (Map and Diagram for all hit reactions)


RXL ----- Hit Reaction Long (Map, Diagram, Summary for all hit reactions)

RXS ----- Hit Reaction Summaries (Map and Summary for all hit reactions)


SPATH ----- Reaction Map and Reaction Diagram for the "short path". Displays all single step reactions which contain a hit substance. Also displays those multistep reactions that have a hit substance in both the first and last steps of the reaction, except for those hit reactions whose steps are totally included within another hit reaction which is displayed

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of combinations include: D TI; D BIB RX; D TI, AU, FCRD. The information is displayed in the same order as the specification. All of the formats, except CRD, CRDREF, FHIT, PATH, FPATH, SPATH, FSPATH, FCRD, FCRDREF, HIT, RX, RXG, RXS, SCAN, and OCC, may be used with the DISPLAY command to display the record for a specified Accession Number.

L3 ANSWER 1 OF 10 CASREACT COPYRIGHT 2008 ACS on STN



1. FeCl<sub>3</sub>, N<sub>2</sub>H<sub>4</sub>, Water,  
MeOH  
2. HCO<sub>2</sub>H, HCl, Water →



81%

REF: Heterocycles, 71(1), 39-48; 2007

NOTE: key intermediate

CON: STAGE(1) 1 hour, reflux; 2 hours, reflux



STAGE(2) room temperature -> 130 deg C; 3 hours, 130 deg C

=> d cbib abs crd 1-10


L3 ANSWER 1 OF 10 CASREACT COPYRIGHT 2008 ACS on STN

146:421938 One-pot conversion of 2-nitrobenzonitriles to quinazolin-4(3H)-ones and synthesis of gefitinib and erlotinib hydrochloride. Chandregowda, Venkateshappa; Rao, Gudapati Venkateswara; Reddy, Goukanapalli Chandrasekara (Vittal Mallya Scientific Research Foundation, Bangalore, 560004, India). Heterocycles, 71(1), 39-48 (English) 2007. CODEN: HTCYAM. ISSN: 0385-5414. Publisher: Japan Institute of Heterocyclic Chemistry.


GI



AB A simple and efficient one-pot conversion of 2-nitrobenzonitriles to quinazolin-4(3H)-ones involving reduction, formylation, hydrolysis and cyclization is reported. E.g., quinazolin-4(3H)-one derivative I was prepared with 85% yield by reacting the corresponding 2-nitrobenzonitrile II with hydrazine using  $\text{FeCl}_3$  in  $\text{MeOH}/\text{H}_2\text{O}$  followed by treating the reaction mixture with formic acid and  $\text{HCl}$ . These quinazolinones have been used for making in economical way the anticancer drug mols. gefitinib (Iressa) and erlotinib  $\text{HCl}$  (Tarceva).



1.  $\text{FeCl}_3$ ,  $\text{N}_2\text{H}_4$ , Water,  
 $\text{MeOH}$   
 2.  $\text{HCO}_2\text{H}$ ,  $\text{HCl}$ , Water  $\rightarrow$




NOTE: key intermediate


CON: STAGE(1) 1 hour, reflux; 2 hours, reflux

STAGE(2) room temperature  $\rightarrow$  130 deg C; 3 hours, 130 deg C

RX(37) OF 57 - 2 STEPS



- 1.1. HNO<sub>3</sub>, Water, Ac<sub>2</sub>O
- 1.2. NH<sub>4</sub>OH, Water
- 2.1. FeCl<sub>3</sub>, N<sub>2</sub>H<sub>4</sub>,  
Water, MeOH, →
- 2.2. HCO<sub>2</sub>H, HCl, Water



81%

NOTE: 1) 75% overall yield from 3,4-dihydroxybenzaldehyde, regioselective, 2) key intermediate



CON: STEP(1.1) 45 - 50 deg C; 8 hours, 45 - 50 deg C

STEP(1.2) pH 8

STEP(2.1) 1 hour, reflux; 2 hours, reflux

STEP(2.2) room temperature -> 130 deg C; 3 hours, 130 deg C

RX(48) OF 57 - 3 STEPS



81%

NOTE: 2) 75% overall yield from 3,4-dihydroxybenzaldehyde, regioselective, 3) key intermediate

CON: STEP(1.1) 25 deg C -> reflux; 1 hour, reflux

STEP(1.2) room temperature -> 110 deg C; 3 hours, 110 deg C

STEP(1.3) pH 8


STEP(2.1) 45 - 50 deg C; 8 hours, 45 - 50 deg C

STEP(2.2) pH 8

STEP(3.1) 1 hour, reflux; 2 hours, reflux

STEP(3.2) room temperature -> 130 deg C; 3 hours, 130 deg C

RX(49) OF 57 - 4 STEPS



NOTE: 3) 75% overall yield from 3,4-dihydroxybenzaldehyde, regioselective, 4) key intermediate


CON: STEP(1.1) room temperature -> 100 deg C; 3 hours, 100 deg C  
 STEP(2.1) 25 deg C -> reflux; 1 hour, reflux  
 STEP(2.2) room temperature -> 110 deg C; 3 hours, 110 deg C  
 STEP(2.3) pH 8  
 STEP(3.1) 45 - 50 deg C; 8 hours, 45 - 50 deg C  
 STEP(3.2) pH 8  
 STEP(4.1) 1 hour, reflux; 2 hours, reflux  
 STEP(4.2) room temperature -> 130 deg C; 3 hours, 130 deg C

L3 ANSWER 2 OF 10 CASREACT COPYRIGHT 2008 ACS on STN

146:274315 Improved synthesis of substituted 6,7-dihydroxy-4-quinazolineamines: tandutinib, erlotinib and gefitinib. Knesl, Petr; Roeseling, Dirk; Jordis, Ulrich (Institute of Applied Synthetic Chemistry, Vienna University of Technology, Vienna, 1060, Austria). Molecules, 11(4), 286-297 (English) 2006. CODEN: MOLEFW. ISSN: 1420-3049. URL: <http://www.mdpi.org/molecules/papers/11040286.pdf> Publisher: Molecular Diversity Preservation International.

AB The synthesis of three substituted 6,7-dihydroxy-4-quinazolineamines: tandutinib, erlotinib and gefitinib in improved yields is reported. The intermediates were characterized by NMR and the purities determined by HPLC.

RX(12) OF 81



CON: 12 hours, 165 - 170 deg C

CCOC(=O)c1cc(ccc1OCCOC)OCCOCCOCOCOc1ccc2c(c1)c(=O)[nH]cnc2=O


CON: STEP(1) room temperature, 50 psi  
STEP(2) 12 hours, 165 - 170 deg C

CCOC(=O)c1ccc(OCCCOC)c(OCCOC)c1COCOCc1cc2c(c1)c(=O)[nH]cnc2C(=O)OCCOC


CON: STEP(1.1) 30 minutes, 0 - 5 deg C; 24 hours, room temperature  
STEP(2) room temperature, 50 psi  
STEP(3) 12 hours, 165 - 170 deg C



RX(54) OF 81 - 4 STEPS



- 1.1. Bu<sub>4</sub>N.I, K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CO
- 1.2. MeOCH<sub>2</sub>CH<sub>2</sub>Cl
2. HNO<sub>3</sub>, Water, AcOH
3. PtO<sub>2</sub>, H<sub>2</sub>, MeOH
4. Formamide



84%

NOTE: 1) alternative preparation shown

CON: STEP(1.1) 20 minutes, room temperature

STEP(1.2) 5 days, reflux

STEP(2.1) 30 minutes, 0 - 5 deg C; 24 hours, room temperature

STEP(3) room temperature, 50 psi


STEP(4) 12 hours, 165 - 170 deg C

L3 ANSWER 3 OF 10 CASREACT COPYRIGHT 2008 ACS on STN

145:271722 Fluorine-18 labeling of 6,7-disubstituted anilinoquinazoline derivatives for positron emission tomography (PET) imaging of tyrosine kinase receptors: Synthesis of 18F-Iressa and related molecular probes. Seimille, Yann; Phelps, Michael E.; Czernin, Johannes; Silverman, Daniel H. S. (Ahmanson Biological Imaging Division, Department of Molecular and Medical Pharmacology, University of California, Los Angeles, CA, 90095-6942, USA). Journal of Labelled Compounds & Radiopharmaceuticals, 48(11), 829-843 (English) 2005. CODEN: JLCRD4. ISSN: 0362-4803. Publisher: John Wiley & Sons Ltd..

AB Inhibitors of tyrosine kinase enzymic activity are a promising new class of antineoplastic agents. Although clin. studies performed over the last decade give more insight on the potential therapeutic applications of such drugs, identification of the individual patients who might benefit from them remains a major challenge. The authors have developed a synthetic strategy for the production of a wide variety of radiolabeled 6,7-disubstituted 4-anilinoquinazolines suitable for non-invasive imaging of tyrosine kinase receptors to predict therapy effectiveness. Three new F-18 labeled radiopharmaceuticals based on the therapeutic agents Tarceva, Iressa, and ZD6474 were synthesized. Decay-corrected yields were 25-40% for a total synthesis time of 120 min, thus providing F-18 labeled tyrosine kinase inhibitors in quantities and times practical for use as PET radiopharmaceuticals.


RX(23) OF 163



62%

NOTE: Niementowski condensation  
CON: 3 hours, 160 deg C


RX(45) OF 163 - 2 STEPS




62%

NOTE: 2) Niementowski condensation  
CON: STEP(1) room temperature  
STEP(2) 3 hours, 160 deg C

RX(74) OF 163 - 3 STEPS



1. HNO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, F<sub>3</sub>CCO<sub>2</sub>H
2. Pd, H<sub>2</sub>, MeOH
3. Formamide, Ammonium formate



62%


NOTE: 3) Niementowski condensation

CON: STEP(1.1) 15 minutes, 0 - 5 deg C; 2 hours, room temperature


STEP(2) room temperature

STEP(3) 3 hours, 160 deg C

RX(75) OF 163 - 4 STEPS



1. MeOCH<sub>2</sub>CH<sub>2</sub>Br, K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CO
2. HNO<sub>3</sub>, CH<sub>2</sub>Cl<sub>2</sub>, F<sub>3</sub>CCO<sub>2</sub>H
3. Pd, H<sub>2</sub>, MeOH
4. Formamide, Ammonium formate



62%

NOTE: 4) Niementowski condensation

CON: STEP(1) 72 hours, reflux

STEP(2.1) 15 minutes, 0 - 5 deg C; 2 hours, room temperature

STEP(3) room temperature


STEP(4) 3 hours, 160 deg C

L3 ANSWER 4 OF 10 CASREACT COPYRIGHT 2008 ACS on STN


145:249220 Preparation of 4-(indol-3-yl)quinazolines as epidermal growth factor receptor inhibitors. Loewe, Werner; Lueth, Anja (Freie Universitaet Berlin, Germany). PCT Int. Appl. WO 2006084882 A2 20060817, 23pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA; RW: AT,

BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IS, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (German). CODEN: PIXXD2. APPLICATION: WO 2006-EP50813 20060209. PRIORITY: DE 2005-102005007151 20050211.

GI





I



II

AB Title compds. I [R1 = methoxy, ethoxy, propoxy, etc.; R2 = H, methoxy, ethoxy, etc.; R3 = (R3')<sub>n</sub>; R3' = halo, halo substituted benzyloxy, alkyl, etc.; n = 1-3] and their pharmaceutically acceptable salts were prepared. For example, Mg mediated coupling of 4-chloro-6,7-dimethoxyquinazoline and 5-bromoindole afforded quinazoline II. In epidermal growth factor receptor inhibition assays, 4-examples of compds. I exhibited 61-81% inhibition at 0.1  $\mu$ M.

RX(5) OF 18



CON: 24 hours, reflux


L3 ANSWER 5 OF 10 CASREACT COPYRIGHT 2008 ACS on STN

144:370039 Synthesis and biological evaluation of allenic quinazolines using palladium-catalyzed hydride-transfer reaction. Nakamura, Hiroyuki; Onagi, Shinya (Department of Chemistry, Faculty of Science, Gakushuin University, Mejiro, Tokyo, 171-8588, Japan). Tetrahedron Letters, 47(15), 2539-2542 (English) 2006. CODEN: TELEAY. ISSN: 0040-4039. Publisher: Elsevier B.V..

AB Allenic quinazolines were designed as mimics of Tarceva, which is an epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor, and synthesized from the corresponding 4-(iodoanilino)quinazolines or 4-(iodophenoxy)quinazolines with N,N-dicyclohexylprop-2-ynylamine by the Sonogashira coupling followed by palladium-catalyzed hydride-transfer


reaction. Cell growth inhibition of the new compds. toward A431, Kato III, SKBR3, and HepG2 was examined. One of the compds. synthesized, showed a similar cell growth inhibition to Tarceva. Moreover, two other compds. exhibited a specific growth inhibition toward Kato III cells (IC50 = 12 and 4.7  $\mu\text{M}$ , resp.), although a significant inhibition toward other three cell lines was not observed at a 100  $\mu\text{M}$  concentration of compds.

RX(16) OF 171




CON: 160 deg C

RX(20) OF 171




CON: reflux

RX (21) OF 171






CON: reflux

RX (24) OF 171





CON: 60 deg C




RX(25) OF 171



+




RX(25) OF 171




60%

CON: 60 deg C

RX(32) OF 171




$\xrightarrow[\text{R: 76858-94-1, CHCl}_3]{\text{Pd complex,}}$



CON: 100 deg C


RX(33) OF 171



$\xrightarrow[\text{R: 76858-94-1, CHCl}_3]{\text{Pd complex,}}$




RX(33) OF 171




61%

CON: 100 deg C

RX(51) OF 171 - 2 STEPS






1. Pd, H<sub>2</sub>, EtOH  
2. Formamide



90%

CON: STEP(2) 160 deg C




RX(55) OF 171 - 2 STEPS



61%

CON: STEP(1) reflux  
 STEP(2) reflux


RX(56) OF 171 - 2 STEPS



99%

CON: STEP(1) reflux  
 STEP(2) reflux


RX(59) OF 171 - 2 STEPS




24%

CON: STEP(1) reflux  
STEP(2) 60 deg C

RX(60) OF 171 - 2 STEPS




RX(60) OF 171 - 2 STEPS




60%

CON: STEP(1) reflux  
STEP(2) 60 deg C

RX(63) OF 171 - 2 STEPS




1. Pd(PPh<sub>3</sub>)<sub>4</sub>, CuI,  
Et<sub>3</sub>N, MeCN  
2. Pd complex,  
R: 76858-94-1,  
CHCl<sub>3</sub>



55%


CON: STEP(1) 60 deg C  
STEP(2) 100 deg C

RX(64) OF 171 - 2 STEPS




CON: STEP(1) 60 deg C  
 STEP(2) 100 deg C

RX(89) OF 171 - 3 STEPS




CON: STEP(3) 160 deg C

RX(90) OF 171 - 4 STEPS




1.  $\text{MeOCH}_2\text{CH}_2\text{Br}$ ,  $\text{K}_2\text{CO}_3$ ,  
 $\text{Me}_2\text{CO}$ ,  $\text{Bu}_4\text{N}^+\text{I}^-$
2.  $\text{HNO}_3$ , Water,  $\text{Ac}_2\text{O}$
3.  $\text{Pd}$ ,  $\text{H}_2$ ,  $\text{EtOH}$
4. Formamide




90%

CON: STEP(1) reflux  
STEP(4) 160 deg C

RX(95) OF 171 - 3 STEPS




1. Formamide
2.  $\text{EtN}(\text{Pr}-i)_2$ ,  $\text{POCl}_3$ ,  
 $\text{PhMe}$
3.  $\text{K}_2\text{CO}_3$ ,  $\text{Me}_2\text{CHOH}$




61%

CON: STEP(1) 160 deg C  
STEP(2) reflux  
STEP(3) reflux

RX(96) OF 171 - 3 STEPS




1. Formamide
2.  $\text{EtN}(\text{Pr-i})_2$ ,  $\text{POCl}_3$ ,  
PhMe
3.  $\text{K}_2\text{CO}_3$ ,  $\text{Me}_2\text{CHOH}$




99%

CON: STEP(1) 160 deg C  
 STEP(2) reflux  
 STEP(3) reflux

RX(99) OF 171 - 4 STEPS




1. Pd,  $\text{H}_2$ , EtOH
2. Formamide
3.  $\text{EtN}(\text{Pr-i})_2$ ,  $\text{POCl}_3$ ,  
PhMe
4.  $\text{K}_2\text{CO}_3$ ,  $\text{Me}_2\text{CHOH}$




61%

CON: STEP(2) 160 deg C  
 STEP(3) reflux  
 STEP(4) reflux

RX(100) OF 171 - 4 STEPS




1. Pd, H<sub>2</sub>, EtOH
2. Formamide
3. EtN(Pr-i)<sub>2</sub>, POCl<sub>3</sub>,  
PhMe
4. K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CHOH




99%

CON: STEP(2) 160 deg C  
 STEP(3) reflux  
 STEP(4) reflux

RX(103) OF 171 - 3 STEPS



1. EtN(Pr-i)<sub>2</sub>, POCl<sub>3</sub>,  
PhMe
2. K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CHOH
3. Pd(PPh<sub>3</sub>)<sub>4</sub>, CuI,  
Et<sub>3</sub>N, MeCN





24%

CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(3) 60 deg C




RX(104) OF 171 - 3 STEPS



1. EtN(Pr-i)2, POC13, PhMe
2. K2CO3, Me2CHOH
3. Pd(PPh3)4, CuI, Et3N, MeCN




RX(104) OF 171 - 3 STEPS



60%


CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(3) 60 deg C

RX(107) OF 171 - 4 STEPS



1. Formamide
2. EtN(Pr-i)2, POC13, PhMe
3. K2CO3, Me2CHOH
4. Pd(PPh3)4, CuI, Et3N, MeCN



RX(107) OF 171 - 4 STEPS



24%


CON: STEP(1) 160 deg C  
STEP(2) reflux  
STEP(3) reflux  
STEP(4) 60 deg C

RX(108) OF 171 - 4 STEPS



1. Formamide
2.  $\text{EtN}(\text{Pr-i})_2$ ,  $\text{POCl}_3$ ,  $\text{PhMe}$
3.  $\text{K}_2\text{CO}_3$ ,  $\text{Me}_2\text{CHOH}$
4.  $\text{Pd}(\text{PPh}_3)_4$ ,  $\text{CuI}$ ,  $\text{Et}_3\text{N}$ ,  $\text{MeCN}$


RX(108) OF 171 - 4 STEPS




60%

CON: STEP(1) 160 deg C  
STEP(2) reflux  
STEP(3) reflux  
STEP(4) 60 deg C

RX(113) OF 171 - 3 STEPS




1. K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CHOH  
2. Pd(PPh<sub>3</sub>)<sub>4</sub>, CuI,  
Et<sub>3</sub>N, MeCN  
3. Pd complex,  
R: 76858-94-1,  
CHCl<sub>3</sub>



55%


CON: STEP(1) reflux  
STEP(2) 60 deg C  
STEP(3) 100 deg C

RX(114) OF 171 - 4 STEPS




CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(3) 60 deg C  
 STEP(4) 100 deg C

RX(115) OF 171 - 3 STEPS




CON: STEP(1) reflux  
 STEP(2) 60 deg C  
 STEP(3) 100 deg C

RX(116) OF 171 - 4 STEPS




CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(3) 60 deg C  
 STEP(4) 100 deg C

RX(121) OF 171 - 5 STEPS




CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(5) 160 deg C

RX(126) OF 171 - 5 STEPS




1. HNO<sub>3</sub>, Water, Ac<sub>2</sub>O
2. Pd, H<sub>2</sub>, EtOH
3. Formamide
4. EtN(Pr-i)<sub>2</sub>, POCl<sub>3</sub>, PhMe
5. K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CHOH




CON: STEP(3) 160 deg C  
STEP(4) reflux  
STEP(5) reflux

RX(127) OF 171 - 5 STEPS




1. HNO<sub>3</sub>, Water, Ac<sub>2</sub>O
2. Pd, H<sub>2</sub>, EtOH
3. Formamide
4. EtN(Pr-i)<sub>2</sub>, POCl<sub>3</sub>, PhMe
5. K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CHOH




CON: STEP(3) 160 deg C  
STEP(4) reflux  
STEP(5) reflux

RX(130) OF 171 - 6 STEPS




1. MeOCH<sub>2</sub>CH<sub>2</sub>Br  
4. Formamide




61%

CON: STEP(1) reflux  
STEP(4) 160 deg C  
STEP(5) reflux  
STEP(6) reflux

RX(131) OF 171 - 6 STEPS




1. MeOCH<sub>2</sub>CH<sub>2</sub>Br  
4. Formamide




99%

CON: STEP(1) reflux  
STEP(4) 160 deg C  
STEP(5) reflux  
STEP(6) reflux

RX(134) OF 171 - 7 STEPS




1. MeOH  
2. MeOCH<sub>2</sub>CH<sub>2</sub>Br  
5. Formamide




61%

CON: STEP(1) reflux  
STEP(2) reflux  
STEP(5) 160 deg C  
STEP(6) reflux  
STEP(7) reflux

RX(135) OF 171 - 7 STEPS



1. MeOH  
2. MeOCH<sub>2</sub>CH<sub>2</sub>Br  
5. Formamide





99%

CON: STEP(1) reflux  
STEP(2) reflux  
STEP(5) 160 deg C  
STEP(6) reflux  
STEP(7) reflux




RX(138) OF 171 - 5 STEPS





1. Pd, H<sub>2</sub>, EtOH
2. Formamide
3. EtN(Pr-i)<sub>2</sub>, POCl<sub>3</sub>, PhMe
4. K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CHOH
5. Pd(PPh<sub>3</sub>)<sub>4</sub>, CuI, Et<sub>3</sub>N, MeCN

RX(138) OF 171 - 5 STEPS




CON: STEP(2) 160 deg C  
 STEP(3) reflux  
 STEP(4) reflux  
 STEP(5) 60 deg C

RX(139) OF 171 - 5 STEPS



1. Pd, H<sub>2</sub>, EtOH
2. Formamide
3. EtN(Pr-i)<sub>2</sub>, POCl<sub>3</sub>, PhMe
4. K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CHOH
5. Pd(PPh<sub>3</sub>)<sub>4</sub>, CuI, Et<sub>3</sub>N, MeCN


RX(139) OF 171 - 5 STEPS



60%

CON: STEP(2) 160 deg C  
 STEP(3) reflux  
 STEP(4) reflux  
 STEP(5) 60 deg C


RX(142) OF 171 - 6 STEPS




24%

CON: STEP(3) 160 deg C  
 STEP(4) reflux  
 STEP(5) reflux  
 STEP(6) 60 deg C

RX(143) OF 171 - 6 STEPS




RX(143) OF 171 - 6 STEPS



60%


CON: STEP(3) 160 deg C  
 STEP(4) reflux  
 STEP(5) reflux  
 STEP(6) 60 deg C

RX(146) OF 171 - 7 STEPS




CON: STEP(1) reflux  
 STEP(4) 160 deg C  
 STEP(5) reflux  
 STEP(6) reflux  
 STEP(7) 60 deg C

RX(147) OF 171 - 7 STEPS




CON: STEP(1) reflux  
 STEP(4) 160 deg C  
 STEP(5) reflux  
 STEP(6) reflux  
 STEP(7) 60 deg C

RX(150) OF 171 - 8 STEPS




CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(5) 160 deg C  
 STEP(6) reflux  
 STEP(7) reflux  
 STEP(8) 60 deg C

RX(151) OF 171 - 8 STEPS




CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(5) 160 deg C  
 STEP(6) reflux  
 STEP(7) reflux  
 STEP(8) 60 deg C

RX(154) OF 171 - 5 STEPS





RX(154) OF 171 - 5 STEPS



CON: STEP(1) 160 deg C  
 STEP(2) reflux  
 STEP(3) reflux  
 STEP(4) 60 deg C  
 STEP(5) 100 deg C


RX(155) OF 171 - 5 STEPS



1. Formamide
2. EtN(Pr-i)2, POC13, PhMe
3. K2CO3, Me2CHOH
4. Pd(PPh3)4, CuI, Et3N, MeCN
5. Pd complex, R:76858-94-1, CHCl3




RX(155) OF 171 - 5 STEPS




61%

CON: STEP(1) 160 deg C  
STEP(2) reflux  
STEP(3) reflux  
STEP(4) 60 deg C  
STEP(5) 100 deg C

RX(158) OF 171 - 6 STEPS




(step 4)



(step 5)


2. Formamide →



55%


CON: STEP(2) 160 deg C  
STEP(3) reflux  
STEP(4) reflux  
STEP(5) 60 deg C  
STEP(6) 100 deg C

RX(159) OF 171 - 6 STEPS



2. Formamide →


RX(159) OF 171 - 6 STEPS



61%


CON: STEP(2) 160 deg C  
 STEP(3) reflux  
 STEP(4) reflux  
 STEP(5) 60 deg C  
 STEP(6) 100 deg C

RX(162) OF 171 - 7 STEPS




CON: STEP(3) 160 deg C  
 STEP(4) reflux  
 STEP(5) reflux  
 STEP(6) 60 deg C  
 STEP(7) 100 deg C

RX(163) OF 171 - 7 STEPS




RX(163) OF 171 - 7 STEPS



61%


CON: STEP(3) 160 deg C  
 STEP(4) reflux  
 STEP(5) reflux  
 STEP(6) 60 deg C  
 STEP(7) 100 deg C

RX(166) OF 171 - 8 STEPS




CON: STEP(1) reflux  
 STEP(4) 160 deg C  
 STEP(5) reflux  
 STEP(6) reflux  
 STEP(7) 60 deg C  
 STEP(8) 100 deg C

RX(167) OF 171 - 8 STEPS




CON: STEP(1) reflux  
 STEP(4) 160 deg C  
 STEP(5) reflux  
 STEP(6) reflux  
 STEP(7) 60 deg C  
 STEP(8) 100 deg C

RX(170) OF 171 - 9 STEPS



CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(5) 160 deg C  
 STEP(6) reflux  
 STEP(7) reflux  
 STEP(8) 60 deg C  
 STEP(9) 100 deg C

RX(171) OF 171 - 9 STEPS



CON: STEP(1) reflux  
 STEP(2) reflux  
 STEP(5) 160 deg C  
 STEP(6) reflux  
 STEP(7) reflux  
 STEP(8) 60 deg C  
 STEP(9) 100 deg C




L3 ANSWER 6 OF 10 CASREACT COPYRIGHT 2008 ACS on STN

142:240390 Improved, high yield synthesis of 3H-quinazolin-4-ones, the key intermediates of recently developed drugs. Oerfi, Laszlo; Waczek, Frigyes; Pato, Janos; Varga, Istvan; Hegymegi-Barakonyi, Balint; Houghten, Richard A.; Keri, Gyoergy (Department of Pharmaceutical Chemistry, Semmelweis University, Budapest, Hung.). Current Medicinal Chemistry, 11(19), 2549-2553 (English) 2004. CODEN: CMCHE7. ISSN: 0929-8673. Publisher: Bentham Science Publishers Ltd..

AB Purine bases and their bioisosteric analogs are widely used as building blocks in combinatorial chemical Recently a great number of fused pyrimidine derivs. became known as potential drug mols. against various types of proliferative diseases, caused by over-expression of protein kinases. One of the most important compound families are quinazolines: e.g. the best inhibitor of EGFR tyrosine kinase is PD153035 (6,7-dimethoxy-4-(3'-bromophenyl)amino-quinazoline) [2] and IRESSA (gefitinib, ZD1839) [3], developed from this compound family, is presently the only one approved and

granted drug by the FDA for the treatment of advanced non-small-cell lung cancer (NSCLC). KF31327 (3-ethyl-8-[2-(4-hydroxymethylpiperidino)benzylamino]-2,3-dihydro-1H-imidazo[4,5-g]-quinazoline-2-thione dihydrochloride) from this group, showed significantly higher inhibitory activity on cyclic GMP-specific phosphodiesterase compared with those of sildenafil (Viagra). The synthetic procedures of the example compds. are based on imidoyl chloride intermediates that were prepared from the appropriate 3H-quinazoline-4-ones. Although the key intermediates, quinazoline-4-ones, have been known since more than hundred years, their synthetic procedures have been improved much only in the past ten years. In this paper we reviewed the efficient synthetic methods of quinazolin-4-ones; and presented a novel, reliable method for their synthesis. There was no considerable effect of microwave-, or traditional thermal activation on the yield and compound purity.

RX(27) OF 30



86%

NOTE: workup: hot aq. NaOH /charcoal decolorization and acidification  
(purity 87%)




CON: 2 hours, 160 deg C

L3 ANSWER 7 OF 10 CASREACT COPYRIGHT 2008 ACS on STN

142:219298 Process for preparation of 6,7-bis(2-methoxyethoxy)quinazolin-4-one. Nishino, Shigeyoshi; Hirotsu, Kenji; Shima, Hidetaka; Oda, Hiroyuki; Suzuki, Shinobu (Ube Industries, Ltd., Japan). PCT Int. Appl. WO 2005012264 A1 20050210, 18 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (Japanese). CODEN: PIXXD2. APPLICATION: WO 2004-JP10965 20040730. PRIORITY: JP 2003-282696 20030730.

AB This invention pertains to a method for producing 6,7-bis(2-methoxyethoxy)quinazolin-4-one, which comprises reacting Et 2-amino-4,5-bis(2-methoxyethoxy)benzoate with a formic acid compound in the presence of an ammonium carboxylate. For example, Et 2-amino-4,5-bis(2-methoxyethoxy)benzoate (preparation given) was reacted with Me orthoformate in MeOH in the presence of NH4OAc to give the title compound (91%). This invention provides a convenient method to prepare the title compound from Et 2-amino-4,5-bis(2-methoxyethoxy)benzoate in high yield.




RX(4) OF 10



91%

CON: 7 hours, 60 - 70 deg C

RX(7) OF 10 - 2 STEPS




91%


CON: STEP(1) 6 hours, 50 - 60 deg C  
STEP(2) 7 hours, 60 - 70 deg C



RX(9) OF 10 - 3 STEPS




1. H2SO4, HNO3
2. Pt, H2, MeOH
3. NH4OAc, CH(OMe)3, MeOH




91%

CON: STEP(1.1) room temperature -> 70 deg C; 2 hours, 70 deg C  
 STEP(2) 6 hours, 50 - 60 deg C  
 STEP(3) 7 hours, 60 - 70 deg C

RX(10) OF 10 - 4 STEPS



1. MeOCH2CH2Cl, K2CO3, DMF
2. H2SO4, HNO3
3. Pt, H2, MeOH
4. NH4OAc, CH(OMe)3, MeOH



91%


CON: STEP(1) 9 hours, 90 - 100 deg C  
 STEP(2.1) room temperature -> 70 deg C; 2 hours, 70 deg C  
 STEP(3) 6 hours, 50 - 60 deg C  
 STEP(4) 7 hours, 60 - 70 deg C

L3 ANSWER 8 OF 10 CASREACT COPYRIGHT 2008 ACS on STN


139:164805 Process for producing 3,4-dihydroquinazolin-4-one derivatives.  
 Nishino, Shigeyoshi; Hirotsu, Kenji; Shima, Hidetaka; Harada, Takashi;  
 Oda, Hiroyuki; Takahashi, Takeshi; Suzuki, Shinobu (Ube Industries, Ltd.,  
 Japan). PCT Int. Appl. WO 2003064399 A1 20030807, 101 pp. DESIGNATED  
 STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH,  
 CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM,  
 HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU,  
 LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD,  
 SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,  
 ZM, ZW; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR,  
 GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR.

(Japanese). CODEN: PIXXD2. APPLICATION: WO 2003-JP805 20030128.  
 PRIORITY: JP 2002-17957 20020128; JP 2002-40929 20020219; JP 2002-82607  
 20020325; JP 2002-168443 20020610; JP 2002-178661 20020619; JP 2002-246657  
 20020827; JP 2002-326752 20021111; JP 2002-349456 20021201.

GI






I



II

AB Disclosed is a process for producing a quinazolin-4-one derivative represented by the following formula (I) (wherein R1, R2, R3, and R4 each represents a group not participating in the reaction and R1, R2, R3, and R4 may be bonded to each other to form a ring) which comprises reacting an anthranilic acid derivative represented by the following formula (II) (wherein R1-R4 are defined as above; R5 represents hydrogen or a hydrocarbon group) with a formic acid derivative (in particular orthoformate ester) in the presence of an ammonium carboxylate. This process gives 3,4-dihydroquinazolin-4-one derivs. I, which are useful as intermediates for drugs and agrochems., from anthranilic acid derivs. in high yields in a simple method under mild conditions and is industrially suitable. Thus, 5-methoxy-4-(3-chloropropoxy)anthranilic acid Me ester 161.5, Me orthoformate 156.5, ammonium acetate 113.7 g, and 300 mL MeOH were added to a 1,000 mL stainless steel pressure vessel, allowed to react at 90-95° and 0.1-0.3 MPa for 8 h, treated with 600 mL H2O, stirred at 0-10° for 1 h, and filtered to give, after washing the crystals with 600 mL H2O and drying them at 60° under reduced pressure, 94% 6-methoxy-7-(3-chloropropoxy)quinazolin-4-one (152.8 g).


RX(11) OF 112



83%

NOTE: high pressure cyclocondensation in a stainless pressure vessel  
 CON: 8 hours, 95 deg C

RX(12) OF 112




NOTE: high pressure cyclocondensation in a stainless pressure vessel  
CON: 6 hours, 110 deg C

L3 ANSWER 9 OF 10 CASREACT COPYRIGHT 2008 ACS on STN


139:164804 Preparation of quinazolinone derivatives and 4,5-substituted-2-formylaminobenzamides as intermediates for said quinazolinones. Shirai, Masashi; Furuya, Toshio (Ube Industries, Ltd., Japan). PCT Int. Appl. WO 2003064377 A1 20030807, 30 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (Japanese). CODEN: PIXXD2. APPLICATION: WO 2003-JP562 20030122. PRIORITY: JP 2002-19583 20020129; JP 2002-77880 20020320.

GI




AB 4,5-Substituted-2-formylaminobenzamides (I) are produced by reacting the corresponding aminobenzamides with formic acid in an organic solvent, and I are further converted into 6,7-substituted-quinazolin-4-ones of the general formula II [R1, R2 = H, (un)substituted alkyl, etc.] by cyclization in the presence of a base. I are pharmaceutical intermediates. 6,7-Dimethoxy-3H-quinazolin-4-one was prepared in 93% yield by the above-mentioned process.

RX(7) OF 12




1. NaOH, Water  
2. HCl, Water




84%

CON: STAGE(1) 30 minutes, 25 deg C, pH 13.5  
STAGE(2) pH 7.6

RX(11) OF 12 - 2 STEPS



1. HCO<sub>2</sub>H  
2.1. NaOH, Water  
2.2. HCl, Water



84%


CON: STEP(1.1) room temperature -> 5 deg C; 5 deg C -> 25 deg C;  
48 hours, 25 deg C  
STEP(2.1) 30 minutes, 25 deg C, pH 13.5  
STEP(2.2) pH 7.6

L3 ANSWER 10 OF 10 CASREACT COPYRIGHT 2008 ACS on STN


137:263058 Preparation of quinazoline derivative as intermediates for antitumor agent. Wang, Wi-Chi; Iseki, Eiichi; Imamiya, Katsuyuki (Sumika Fine Chemicals Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP 2002293773 A 20021009, 9 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 2001-100183 20010330.

AB 6,7-Bis(2-methoxyethoxy)-4(1H)-quinazolinone, an intermediate for the antitumor agent N-(3-ethylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, is prepared in several steps from Et 3,4-dihydroxybenzoate and 2-methoxyethyl mesylate via cyclocondensation of Et 2-amino-4,5-bis(2-methoxyethoxy)benzoate with ammonium formate.


RX(5) OF 15




Ammonium formate,  
Formamide




RX(9) OF 15 - 2 STEPS




1. Pt, H<sub>2</sub>, MeOH  
2. Ammonium formate,  
Formamide




RX(13) OF 15 - 3 STEPS




1. H<sub>2</sub>SO<sub>4</sub>, HNO<sub>3</sub>, AcOH  
2. Pt, H<sub>2</sub>, MeOH  
3. Ammonium formate,  
Formamide




RX(14) OF 15 - 4 STEPS




1. Bu<sub>4</sub>N.I, K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CO
2. H<sub>2</sub>SO<sub>4</sub>, HNO<sub>3</sub>, AcOH
3. Pt, H<sub>2</sub>, MeOH
4. Ammonium formate, Formamide



RX(15) OF 15 - 5 STEPS



1. MeCH<sub>2</sub>CH<sub>2</sub>OH, N-Methylmorpholine, THF
2. Bu<sub>4</sub>N.I, K<sub>2</sub>CO<sub>3</sub>, Me<sub>2</sub>CO
3. H<sub>2</sub>SO<sub>4</sub>, HNO<sub>3</sub>, AcOH
4. Pt, H<sub>2</sub>, MeOH
5. Ammonium formate, Formamide



=>